



Cobalt charge states in sodium cobaltates at intermediate dopings

G. Lang^{a,*}, J. Bobroff^a, H. Alloul^a, I. Mukhamedshin^{a,b}, G. Collin^c, N. Blanchard^a^a Laboratoire de Physique des Solides, UMR8502, University Paris-Sud, 91405 Orsay, France^b Physics Department, Kazan State University, 420008 Kazan, Russia^c Laboratoire Léon Brillouin, CE Saclay, CEA-CNRS, 91191 Gif-sur-Yvette, France

ARTICLE INFO

Article history:

Received 27 June 2008

Received in revised form

18 September 2008

Accepted 19 September 2008

Available online 20 November 2008

PACS:

71.27.+a

76.60.-k

75.20.-g

Keywords:

Transition metal alloys and compounds

Electronic properties

Nuclear resonances

ABSTRACT

A likely crucial parameter in explaining the peculiar magnetism of Na_xCoO_2 cobaltates in their paramagnetic state is their charge and spin states in the cobalt layers, responsible for the electronic properties. We examine here the situation for $0.50 \leq x \leq 0.62$, for which it is shown that an increasing amount of Co^{3+} ions appears among $\sim 3.5+$ ions for increasing x . Together with our previous results, this defines two regions in the phase diagram, with homogeneous Co states for $x \leq 0.5$ and heterogeneous states for $x > 0.5$. The links with the sodium structural ordering and the nature of magnetism are discussed.

© 2008 Elsevier B.V. All rights reserved.

1. Introduction

Sodium cobaltates have recently attracted considerable attention following the discovery of hints of strong correlations [1] and of superconductivity [2]. The latter is remarkable as cobaltates feature a layered structure, reminding one of the high-temperature superconducting cuprates, with layers of edge-sharing CoO_6 octahedra separated by layers of Na ions. On changing the sodium content x , one changes the doping in the Co layers. In a naive picture, going from $x=0$ to $x=1$ results in a progressive replacement of Co^{4+} ions (magnetic, $S=1/2$) with Co^{3+} ions (nonmagnetic, $S=0$). One would thus expect reduced magnetism as x is increased, which is not what is observed experimentally. Indeed, except for the band-insulator limit $x=1$ where we have verified the nonmagnetic $3+$ state of all cobalts [3], magnetic correlations are ubiquitous, with a crossover at $x^* \sim 0.6$ from a high x weakly-ferromagnetic itinerant behavior to a low x antiferromagnetic correlation regime [4,5].

A crucial point to explain this behavior may be the actual cobalt charge and spin states in the system. Previous results by us showed that there is charge disproportionation for $0.67 \leq x \leq 0.75$ [5], while

it is absent for $x \leq 0.5$ [6,7]. Here we focus on the $0.50 \leq x \leq 0.62$ intermediate doping region, which we studied extensively, especially in terms of magnetic correlations in [4]. We show there is heterogeneity of the cobalts states, with increasing amounts of Co^{3+} ions among $\sim 3.5+$ ions for x increasing above 0.5. The latter is thus a threshold for charge disproportionation but does not coincide with the above-mentioned threshold $x^* \sim 0.6$ between correlation regimes, and is argued to be linked to the Na ordering.

2. Results and discussion

To establish the different Co states present for $0.55 \leq x \leq 0.62$, we performed ^{59}Co (spin $I=7/2$) nuclear magnetic resonance, taking advantage of its sensitivity to the local magnetic and charge environment of Co nuclei (see [4] for experimental details). While spectra obtained at low temperature seem to indicate that the $\sim 3.5+$ Co sites detected for $x=0.5$ are still present (not shown), we notice the presence of a new cobalt NMR signal (left panel of Fig. 1): upon increasing x above 0.5, a set of quadrupolar satellites (marked with stars) develops on each side of the central resonance line. The corresponding Co sites have a resonance much less temperature-dependent than the $\sim 3.5+$ sites, with isotropic NMR shift values stable around 2%, and also with a slower transverse relaxation rate. These are the fingerprints of the Co^{3+} site in the $x=1$ and $x=0.67$ phases [3,8], showing that such sites start appearing beyond $x=0.5$.

* Corresponding author. Tel.: +49 351 4659581; fax: +49 351 4659313.

E-mail address: g.m.lang@ifw-dresden.de (G. Lang).